

[Click here to view linked References](#)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65	Noname manuscript No. (will be inserted by the editor)
---	--

Topological properties of function spaces over ordinals

Saak Gabrielyan · Jan Grebík · Jerzy Kąkol · Lyubomyr Zdomskyy

Received: date / Accepted: date

Abstract A topological space X is said to be an *Ascoli space* if any compact subset \mathcal{K} of $C_k(Y)$ is evenly continuous. This definition is motivated by the classical Ascoli theorem. We study the $k_{\mathbb{R}}$ -property and the Ascoli property of $C_p(\kappa)$ and $C_k(\kappa)$ over ordinals κ . We prove that $C_p(\kappa)$ is always an Ascoli space, while $C_p(\kappa)$ is a $k_{\mathbb{R}}$ -space iff the cofinality of κ is countable. In particular, this provides the first C_p -example of an Ascoli space which is not a $k_{\mathbb{R}}$ -space, namely $C_p(\omega_1)$. We show that $C_k(\kappa)$ is Ascoli iff $\text{cf}(\kappa)$ is countable iff $C_k(\kappa)$ is metrizable.

The second author was supported by the GACR project 15-34700L and RVO: 67985840. The third author was supported by Generalitat Valenciana, Conselleria d'Educació, Cultura i Esport, Spain, Grant PROMETEO/2013/058 and by the GAČR project 16-34860L and RVO: 67985840, and gratefully acknowledges also the financial support he received from the Kurt Gödel Research Center in Wien for his research visit in days 15.04-24.04 2016. The fourth author would like to thank the Austrian Science Fund FWF (Grant I 1209-N25) for generous support for this research. The collaboration of the second and the fourth authors was partially supported by the Czech Ministry of Education grant 7AMB15AT035 and RVO: 67985840.

S. Gabrielyan
Department of Mathematics, Ben-Gurion University of the Negev, Beer-Sheva, P.O. 653,
Israel
E-mail: saak@math.bgu.ac.il

J. Grebík
Institute of Mathematics, Czech Academy of Sciences, Czech Republic
E-mail: Greboshrabos@seznam.cz

J. Kąkol
A. Mickiewicz University 61 – 614 Poznań, Poland and Institute of Mathematics, Czech
Academy of Sciences, Czech Republic
E-mail: kakol@amu.edu.pl

L. Zdomskyy
Kurt Gödel Research Center for Mathematical Logic, University of Vienna, A-1090 Wien,
Austria
E-mail: lzdmsky@gmail.com

Keywords $C_p(X) \cdot C_k(X) \cdot$ Ascoli $\cdot k_{\mathbb{R}}$ -space \cdot ordinal space

Mathematics Subject Classification (2010) MSC 54C35 \cdot MSC 54F05 \cdot MSC 46A08 \cdot MSC 54E18

1 Introduction

The study of topological properties of function spaces is quite an active area of research attracting specialists both from topology and functional analysis, see for example [1,3,6,9,13] and references therein. In the following diagram we select the most important compact type properties generalizing metrizable

$$\text{metric} \implies \begin{matrix} \text{Fréchet-} \\ \text{Urysohn} \end{matrix} \implies k\text{-space} \implies k_{\mathbb{R}}\text{-space} \implies \text{Ascoli},$$

and note that none of these implications is reversible, see [3,4] (all relevant definitions are given in the next section).

For a Tychonoff topological space X , we denote by $C_k(X)$ and $C_p(X)$ the space $C(X)$ of all continuous real-valued functions on X endowed with the compact-open topology and the topology of pointwise convergence, respectively.

It is well-known that $C_p(X)$ is metrizable if and only if X is countable. Pytkeev, Gerlitz and Nagy (see §3 of [1]) characterized spaces X for which $C_p(X)$ is Fréchet–Urysohn or a k -space (these properties coincide for spaces of the form $C_p(X)$). The authors in [5] obtained some sufficient conditions on X for which the space $C_p(X)$ is an Ascoli space. Recall that X is called an *Ascoli space* if any compact subset \mathcal{K} of $C_k(X)$ is evenly continuous (or, equivalently, if the natural evaluation map $X \hookrightarrow C_k(C_k(X))$ is an embedding, see [3]).

Every linear order $<$ on a set X generates a natural topology on X whose subbase consists of sets of the form $\{z : z < x\}$ and $\{z : z > x\}$, where $x \in X$. Spaces X whose topology is generated by some linear order are called *linearly ordered topological spaces*. Ordinals with the topology generated by their natural wellorder form an interesting class of linearly ordered topological spaces, and function spaces over them give a good source of (counter)examples in the corresponding theory. For instance, the space $C_p(\omega_1)$ is Lindelöf, see [13]. On the other hand, Arhangel'skii showed in [2] that the space $C_p(\omega_1 + 1)$ is not normal. In [8] Gul'ko proved that there are no two distinct natural number n and m for which the powers $C_p(\omega_1)^n$ and $C_p(\omega_1)^m$ are homeomorphic. In [11] Morris and Wulbert observed that $C_k(\omega_1)$ is not barrelled.

In this short note we provide complete characterizations of those ordinals κ for which $C_p(\kappa)$ and $C_k(\kappa)$ are $k_{\mathbb{R}}$ -spaces or Ascoli spaces. The following theorems are the main results of the paper.

Theorem 1.1 *For every ordinal κ the space $C_p(\kappa)$ is Ascoli.*

Denote by $\text{cf}(\kappa)$ the cofinality of an ordinal κ .

Theorem 1.2 *For an ordinal κ , the space $C_p(\kappa)$ is a $k_{\mathbb{R}}$ -space if and only if $\text{cf}(\kappa) \leq \omega$ if and only if $C_p(\kappa)$ is Fréchet–Urysohn.*

Theorems 1.1 and 1.2 show that the space $C_p(\omega_1)$ is Ascoli but not a $k_{\mathbb{R}}$ -space. This answers Question 6.8 in [6] for spaces $C_p(X)$ in the affirmative, and complements [4, 3.3.E] asserting that for uncountable discrete X the space $C_p(X) = \mathbb{R}^X$ is a $k_{\mathbb{R}}$ -space but not a k -space.

Theorem 1.3 *For an ordinal κ , the space $C_k(\kappa)$ is an Ascoli space if and only if $\text{cf}(\kappa) \leq \omega$, so $C_k(\kappa)$ is complete and metrizable.*

2 Proofs

Below we recall some topological concepts used in Theorems 1.1 and 1.3, for other notions we refer the reader to the book [4]. A k -cover \mathcal{U} of a topological space X is a family of subsets of X such that every compact subset of X is contained in some member of \mathcal{U} .

Definition 2.1 *A topological space X is*

- hemicompact if there exists a countable k -cover of X consisting of compacts;
- realcompact if it can be embedded as a closed subset to \mathbb{R}^λ for some cardinal λ ;
- a $k_{\mathbb{R}}$ -space if a real-valued function f on X is continuous if and only if its restriction $f|_K$ to any compact subset K of X is continuous;
- scattered if every non-empty subspace A of X has an isolated point in A .

Recall that an ordinal κ is *limit* if there is no α such that $\kappa = \alpha + 1$, otherwise κ is called a *successor* ordinal. The *cofinality* $\text{cf}(\kappa)$ of a limit ordinal number κ is the smallest ordinal α which is the order type of a cofinal subset of κ . If κ is a successor ordinal we set $\text{cf}(\kappa) = 1$.

The following simple facts should be well-known (for (i) see [7, § 5.11], the other ones are straightforward).

- Lemma 2.2** (i) κ is compact if and only if it is a successor;
(ii) κ is hemicompact non-countably compact if and only if $\text{cf}(\kappa) = \omega$;
(iii) κ is countably compact non-compact if and only if $\text{cf}(\kappa) > \omega$.

For the convenience of the reader we recall also the following two results.

Proposition 2.3 ([6]) *Assume X admits a family $\mathcal{U} = \{U_i : i \in I\}$ of open subsets of X , a subset $A = \{a_i : i \in I\} \subset X$ and a point $z \in X$ such that: (i) $a_i \in U_i$ for every $i \in I$, (ii) $|\{i \in I : C \cap U_i \neq \emptyset\}| < \infty$ for each compact subset C of X , and (iii) z is a cluster point of A . Then X is not an Ascoli space.*

A family $\{A_i\}_{i \in I}$ of subsets of a set X is said to be *point-finite* if the set $\{i \in I : x \in A_i\}$ is finite for every $x \in X$. A family $\{A_i\}_{i \in I}$ of subsets of a topological space X is called *strongly point-finite* if for every $i \in I$, there exists an open set U_i of X such that $A_i \subseteq U_i$ and $\{U_i\}_{i \in I}$ is point-finite. Following Sakai [12], a topological space X is said to have the *property* (κ) if every pairwise disjoint sequence of finite subsets of X has a strongly point-finite subsequence. The following result is proved in [5].

Theorem 2.4 *If $C_p(X)$ is Ascoli, then X has the property (κ) .*

It is well-known that ordinals are locally compact and scattered (for the last property we note that the smallest element of a subset A of X is isolated in A). The following proposition is of independent interest, it generalizes Corollary 1.5 of [5] and immediately implies Theorem 1.1.

Proposition 2.5 *Let X be a locally compact space. Then $C_p(X)$ is Ascoli if and only if X is scattered.*

Proof The “only if” part follows from Theorem 2.4 combined with the fact that the property (κ) is preserved by subspaces, along with the fact that every compact space with the property (κ) is scattered, see [12, Theorem 3.2]. For the “if” direction consider the one-point compactification $X^* = X \cup \{x_\infty\}$ of X and note that it is scattered. Therefore $C_p(X^*)$ is Fréchet-Urysohn by [1, II.7.16], and hence so is its subspace Z consisting of those continuous $f : X^* \rightarrow \mathbb{R}$ such that $f(x_\infty) = 0$. Now it is easy to see that $Z \upharpoonright X = \{f \upharpoonright X : f \in Z\} \subset C_p(X)$ is homeomorphic to Z and is dense in $C_p(X)$. Therefore $C_p(X)$ has a dense Fréchet-Urysohn subspace, and hence every function f belongs to a dense Ascoli subspace $f + Z$ of $C_p(X)$. Thus $C_p(X)$ is Ascoli by Proposition 5.10 of [3]. \square

Below we prove Theorems 1.2 and 1.3.

Proof of Theorem 1.2. Let $C_p(\kappa)$ be a $k_{\mathbb{R}}$ -space and suppose towards a contradiction that $\text{cf}(\kappa) > \omega$. Then κ is countably compact by Lemma 2.2. Hence the space $C(\kappa)$ endowed with the sup-norm is a Banach space. Therefore the space $C_p(\kappa)$ admits a stronger normed topology and is angelic by [9, Proposition 9.6]. Since every compact subset of $C_p(\kappa)$ is Fréchet-Urysohn, every sequentially continuous function on $C_p(\kappa)$ is continuous. In particular, every sequentially continuous functional on $C_p(\kappa)$ is continuous. So κ is a realcompact space by Theorem 1.1 of [14]. Being realcompact and pseudocompact the space κ is compact by [4, 3.11.1]. Hence κ is a successor ordinal, a contradiction. Thus $\text{cf}(\kappa) \leq \omega$.

Conversely, let $\text{cf}(\kappa) \leq \omega$. Then κ is hemicompact by Lemma 2.2. Thus $C_p(\kappa)$ is Fréchet-Urysohn by [1, II.7.16]. \square

Since each ordinal α is the set of all smaller ordinals, in the following proof we adopt the following perhaps standard notation: For a function f whose domain is an ordinal κ and $\alpha \in \kappa$ we denote by $f(\alpha)$ the value of f at α , and by $f[\alpha]$ the set $\{f(\beta) : \beta < \alpha\}$.

Proof of Theorem 1.3. Suppose for a contradiction that $\text{cf}(\kappa) > \omega$. We shall use Proposition 2.3 and show that $C_k(\kappa)$ is not Ascoli. For every $\alpha < \kappa$ we define $f_\alpha : \kappa \rightarrow [0, 1]$ by $f_\alpha[\alpha + 1] = \{0\}$ and $f_\alpha[\kappa \setminus (\alpha + 1)] = \{1\}$, and set

$$U_\alpha := \{f \in C_k(\kappa) : f(\alpha) < 1/4, f(\alpha + 1) > 3/4\}.$$

To prove that $C_k(\kappa)$ is not Ascoli it is enough to verify the assumptions of Proposition 2.3 for $\{f_\alpha\}_{\alpha < \kappa}$, $\{U_\alpha\}_{\alpha < \kappa}$ and $0 \in C_k(\kappa)$. Clearly, (i) and (iii) hold true. Let us check (ii). Take any compact $C \subseteq C_k(\kappa)$ and assume, contrary to our claim, that there are infinitely many $\alpha < \kappa$ such that $C \cap U_\alpha \neq \emptyset$. Then there exists a strictly increasing sequence $\{\alpha_n\}_{n < \omega}$ such that $C \cap U_{\alpha_n} \neq \emptyset$. Let $\alpha = \lim \alpha_n$. As $\text{cf}(\kappa) > \omega$ we have $\alpha < \kappa$. By the Ascoli theorem used for $\alpha + 1 \subset \kappa$ and $1/2$ we can find a basic neighborhood O_α of α such that $|h(x) - h(y)| < 1/4$ for all $x, y \in O_\alpha$ and $h \in C$. Take n such that $\alpha_n \in O_\alpha$ and fix $h \in C \cap U_{\alpha_n}$. Then

$$\begin{aligned} \frac{1}{4} &> |h(\alpha_n + 1) - h(\alpha_n)| \\ &\geq |f_{\alpha_n}(\alpha_n + 1) - f_{\alpha_n}(\alpha_n)| - |f_{\alpha_n}(\alpha_n + 1) - h(\alpha_n + 1)| - |h(\alpha_n) - f_{\alpha_n}(\alpha_n)| \\ &> 1 - \frac{1}{4} - \frac{1}{4} = \frac{1}{2}, \end{aligned}$$

which is a contradiction. Thus $\text{cf}(\kappa) \leq \omega$.

Conversely, if $\text{cf}(\kappa) \leq \omega$, then κ is a hemicompact locally compact space by Lemma 2.2. Hence $C_k(\kappa)$ is complete metrizable by Corollary 5.2.2 of [10].

□

References

1. Arhangel'skii A. V.: *Topological function spaces*, Math. Appl. **78**, Kluwer Academic Publishers, Dordrecht (1992)
2. Arhangel'skii A. V.: Normality and dense subspaces, Proc. Amer. Math. Soc. **48** 283–291 (2001)
3. Banakh T., Gabrielyan S.: On the C_k -stable closure of the class of (separable) metrizable spaces, Monatshefte Math. **180** 39–64 (2016)
4. Engelking R.: *General topology*, Panstwowe Wydawnictwo Naukowe, Warszawa (1977)
5. Gabrielyan S., Grebík J., Kąkol J., Zdomsky L.: The Ascoli property for function spaces, Topology Appl. **214** 35–50 (2016)
6. Gabrielyan S., Kąkol J., Plebanek G.: The Ascoli property for function spaces and the weak topology on Banach and Fréchet spaces, Studia Math. **233** 119–139 (2016)
7. Gillman L., Jerison M.: *Rings of continuous functions*, Van Nostrand, New York (1960)
8. Gul'ko S. P.: Spaces of continuous functions on ordinals and ultrafilters, Math. Notes **47** 329–334 (1990)
9. Kąkol J., Kubiś W., Lopez-Pellicer M.: *Descriptive Topology in Selected Topics of Functional Analysis*, Developments in Mathematics, Springer, (2011)
10. McCoy R. A., Ntantu I.: *Topological Properties of Spaces of Continuous Functions*, Lecture Notes in Math. **1315** (1988)
11. Morris P. D., Wulbert D. E.: Functional representation of topological algebras, Pacific J. Math. **22** 323–337 (1967)
12. Sakai M.: Two properties of $C_p(X)$ weaker than Fréchet-Urysohn property, Topology Appl. **153** 2795–2804 (2006)
13. Tkachuk V.: *A C_p -theory problem book, special features of function spaces*, Springer, New York (2014)
14. Wilansky A., Mazur spaces, Internat. J. Math. **4** 39–53 (1981)